



2,4-D Ester

Recommended uses

Late Autumn, early Spring, early Winter.

Weeds controlled in : pasture @ 2-4L/ha

Alligator weed, allseed, apple of peru, Australian flax, Australian sedge, bachelors button, Bathurst bur, beggars' ticks, bitter cress, black nightshade, burdock, bur medick, buttercups, Californian poppy, Canadian fleabane, Cape tulip, castor oil plant, catsear, charlock, chicory, clovers, combind, cresses, cudweed, dandelion, doves foot, Fathen, field bindweed, field cress, foxglove, goats rue, greater bindweed, groundsel, hawbit, hawksbeard, hawkweed, hedge mustard, hemp nettle, hoary cress, horehound, marsh cress, Mexican devil, mint weed, monkey musk, nettles, oxtounge, penny cress, piripiri, plantains, poplars, poroporo, ragwort, red dead nettle, redroot, rushes, selfheal, shepherds purse, silver poplar, Spanish heath, tansy, teasel, thistles (Cali, nodding, Scotch, sow, star, variegated, welted, winged), twin cress, vetch, wart cress, white mustard, wild carrot, wild mustard, wild parsnip, wild radish, wild turnip, willow weed, wineberry.

Use lower rates on seedlings

Livestock withholding

Keep off pastures until the plants have died down, or 2 weeks after spraying.

Clovers

May be damaging to clovers especially red clover in Spring and Autumn
Graze area before spraying to reduce clover and expose weeds.